



TITLE:

# Some results on quotient Aubry sets (Viscosity Solutions of Differential Equations and Related Topics)

AUTHOR(S):

FUJITA, Yasuhiro

---

CITATION:

FUJITA, Yasuhiro. Some results on quotient Aubry sets (Viscosity Solutions of Differential Equations and Related Topics).  
数理解析研究所講究録 2009, 1651: 132-136

ISSUE DATE:

2009-05

URL:

<http://hdl.handle.net/2433/140783>

RIGHT:

# Some results on quotient Aubry sets

Yasuhiro FUJITA \*

Department of Mathematics, University of Toyama,  
Toyama 930-8555, Japan.

December 19, 2008

## 1 Introduction

This note consists of partial results of my recent paper [Fu]. In [Fu], we discuss several topics which are not treated here.

Let  $\Omega$  be an open and connected subset of  $\mathbb{R}^n$ , and  $H : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}$  a given function. In this paper, we consider the Hamilton-Jacobi equation

$$(1.1) \quad H(x, Du(x)) = 0 \quad \text{in } \Omega.$$

Let  $\mathcal{A}$  and  $\overline{\mathcal{A}}$  be, respectively, its (projected) Aubry set and the quotient Aubry set. As for their definitions and properties, see Section 2 below. The quotient Aubry set  $\overline{\mathcal{A}}$  plays an essential role to study viscosity solutions of (1.1) (cf. [CI, DS, FU, I, IM]). In particular, several authors provided sufficient conditions in order that  $\overline{\mathcal{A}}$  is totally disconnected (i.e., every connected component consists of a single point in the topology of  $\overline{\mathcal{A}}$ ) [FFR, M1, M2, S].

In this note, we explain a reason why total disconnectedness of  $\overline{\mathcal{A}}$  is important. Let  $\pi(x)$  be the equivalent class of  $\overline{\mathcal{A}}$  containing  $x \in \mathcal{A}$ . We study how  $\pi(x)$  behaves in  $\mathcal{A}$  when  $\overline{\mathcal{A}}$  is total disconnected. We show that a necessary condition in order that  $\overline{\mathcal{A}}$  is totally disconnected is that  $\pi(x) \supset C(x)$  holds for each  $x \in \mathcal{A}$ . Here,  $C(x)$  is the connected component of  $\mathcal{A}$  containing  $x \in \mathcal{A}$ . On the other hand, we show that if  $\mathcal{A}$  is a compact set in  $\Omega$ , then a necessary and sufficient condition in order that  $\overline{\mathcal{A}}$  is totally disconnected is that  $\pi(x) = C(x)$  holds for each  $x \in \mathcal{A}$ .

The state such that  $\pi(x) = C(x)$  for each  $x \in \mathcal{A}$  is preferable, because we can understand and calculate  $\pi(x)$  of this case clearly. Our result shows that if  $\mathcal{A}$  is a compact set in  $\Omega$ , this preferable state occurs when and only when  $\overline{\mathcal{A}}$  is totally disconnected. This is a reason why we propose that totally disconnectedness of  $\overline{\mathcal{A}}$  is important.

The contents of this note are as follows: In Section 2, we provide some preliminaries. In Section 3, we state our results .

---

\*Supported in part by Grant-in-Aid for Scientific Research, No. 18540165, 18204009, JSPS.

## 2 Preliminaries

Let  $B(x, r) = \{y \in \mathbb{R}^n \mid |y - x| \leq r\}$  for  $x \in \mathbb{R}^n$  and  $r > 0$ . We assume:

$$(A1) \quad H \in C(\Omega \times \mathbb{R}^n).$$

$$(A2) \quad H \text{ is coercive, that is, for any compact subset } K \text{ of } \Omega,$$

$$\lim_{r \rightarrow \infty} \inf \{H(x, p) \mid x \in K, p \in \mathbb{R}^n \setminus B(0, r)\} = \infty.$$

$$(A3) \quad \text{For any } x \in \Omega, \text{ the function } p \mapsto H(x, p) \text{ is convex on } \mathbb{R}^n.$$

$$(A4) \quad \text{There is a continuous viscosity subsolution of (1.1).}$$

Let  $\mathcal{S}$  (resp.,  $\mathcal{S}^-$ ) denotes the space of continuous viscosity solutions (resp., viscosity subsolutions) of (1.1). If necessary, we write  $\mathcal{S}(\Omega)$  and  $\mathcal{S}^-(\Omega)$  for  $\mathcal{S}$  and  $\mathcal{S}^-$ , respectively, in order to refer the domain under consideration. Then, (A4) implies that  $\mathcal{S}^-(\Omega) \neq \emptyset$ .

Next, we explain the (projected) Aubry set for the Hamilton-Jacobi equation (1.1). The Aubry set is defined as follows: Define the function  $d : \Omega \times \Omega \mapsto (-\infty, \infty]$  by

$$(2.1) \quad d(x, y) = \sup \{v(x) - v(y) \mid v \in \mathcal{S}^-(\Omega)\}.$$

Then, by [IM, Theorem 1.4 and Proposition 1.6], we have the following:

$$(2.2) \quad d \text{ is locally Lipschitz continuous on } \Omega \times \Omega.$$

$$(2.3) \quad u(x) - u(y) \leq d(x, y) \text{ for all } u \in \mathcal{S}^-(\Omega) \text{ and } x, y \in \Omega.$$

$$(2.4) \quad \text{For all } y \in \Omega, d(\cdot, y) \in \mathcal{S}^-(\Omega) \text{ and } d(\cdot, y) \in \mathcal{S}(\Omega \setminus \{y\}).$$

$$(2.5) \quad \text{For all } x, y, z \in \Omega, d(x, z) \leq d(x, y) + d(y, z) \text{ and } d(x, x) = 0.$$

$$(2.6) \quad d(x, y) = \inf \left\{ \int_0^t L(\gamma(s), \dot{\gamma}(s)) ds \mid t > 0, \gamma \in \mathcal{C}(x, t; y, 0) \right\},$$

where  $\mathcal{C}(x, t; y, 0)$  is the set of all absolutely continuous curves  $\gamma : [0, t] \mapsto \Omega$  satisfying  $(\gamma(t), \gamma(0)) = (x, y)$ , and  $L \in C(\Omega \times \mathbb{R}^n)$  is the convex conjugate of  $H$  defined by

$$(2.7) \quad L(x, \xi) = \sup \{\xi \cdot p - H(x, p) \mid p \in \mathbb{R}^n\} \quad \text{for } (x, \xi) \in \Omega \times \mathbb{R}^n.$$

The Aubry set  $\mathcal{A}$  is defined by

$$(2.8) \quad \mathcal{A} = \{y \in \Omega \mid d(\cdot, y) \in \mathcal{S}(\Omega)\}.$$

In the following, we assume

$$(A5) \quad \mathcal{A} \neq \emptyset.$$

We note that  $\mathcal{A}$  is a closed set in  $\Omega$ , which is due to the stability of the viscosity property under uniform convergence. The assumptions (A1)-(A5) are considered to be natural to discuss the Aubry set for the Hamilton-Jacobi equation (1.1).

Now, we explain an equivalence relation on  $\mathcal{A}$ , which is important to study  $\mathcal{S}(\Omega)$  and  $\mathcal{S}^-(\Omega)$ . By (2.5), we see that the function  $\lambda : \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{R}$  defined by  $\lambda(x, y) = d(x, y) + d(y, x)$  is a pseudo-metric on  $\mathcal{A}$ , i.e., it is non-negative, symmetric, and satisfies the triangle inequality and  $\lambda(x, x) = 0$ ; but the condition  $\lambda(x, y) = 0$  does not necessarily imply  $x = y$ . Let  $x, y \in \mathcal{A}$ . If  $\lambda(x, y) = 0$ , then we write  $x\delta y$ . This relation  $\delta$  is an equivalence relation on  $\mathcal{A}$ . We set

$$(2.9) \quad \begin{aligned} \pi(y) &= \{z \in \mathcal{A} \mid z\delta y\}, \quad y \in \mathcal{A}, \\ \overline{\mathcal{A}} &= \{\pi(y) \mid y \in \mathcal{A}\}. \end{aligned}$$

Then,  $\pi$  is considered as the canonical surjection from  $\mathcal{A}$  to  $\overline{\mathcal{A}}$ , and we see that  $\overline{\mathcal{A}} = \mathcal{A}/\delta$ . Note that we may regard  $\xi \in \overline{\mathcal{A}}$  as a subset of  $\mathcal{A}$  and  $\xi = \pi^{-1}(\{\xi\})$ . Note also that if  $x \in \pi(y)$ , then  $\pi(x) = \pi(y)$ . We define the function  $\overline{\lambda} : \overline{\mathcal{A}} \times \overline{\mathcal{A}} \rightarrow \mathbb{R}$  by

$$(2.10) \quad \overline{\lambda}(\pi(x), \pi(y)) = d(x, y) + d(y, x).$$

The following proposition is well-known.

**Proposition 1.**  *$\overline{\lambda}$  is well defined, and  $(\overline{\mathcal{A}}, \overline{\lambda})$  is a metric space.*

### 3 Results

In this section, we state our results of this note. For their proofs, see [Fu]. Let  $C(x)$  be the connected component of  $\mathcal{A}$  containing  $x$ . In the following, as the topology of  $\overline{\mathcal{A}}$ , we always consider the one induced by the metric  $\overline{\lambda}$ . Note that, by (2.2) and (2.10),  $\pi$  is a continuous mapping from  $\mathcal{A}$  to  $\overline{\mathcal{A}}$ .

**Proposition 2.** *Assume (A1)-(A5). If  $\overline{\mathcal{A}}$  is totally disconnected, then  $\pi(x) \supset C(x)$  for each  $x \in \mathcal{A}$ .*

By Proposition 2, the condition that  $\pi(x) \supset C(x)$  for each  $x \in \mathcal{A}$  is a necessary condition in order that  $\overline{\mathcal{A}}$  is totally disconnected. Next, we consider a sufficient condition in order that  $\overline{\mathcal{A}}$  is totally disconnected. In the following, we assume:

(A6)  $\mathcal{A}$  is a compact set of  $\Omega$ .

We provide a simple consequence of (A6).

**Lemma 1.** *Assume (A1)-(A6). Then,  $\pi(x)$  is a connected set of  $\mathcal{A}$  for each  $x \in \mathcal{A}$ .*

Now, we are in the position to state our sufficient condition in order that  $\overline{\mathcal{A}}$  is totally disconnected.

**Proposition 3.** *Assume that (A1)-(A6). Then,  $\overline{\mathcal{A}}$  is totally disconnected if and only if  $\pi(x) = C(x)$  for each  $x \in \mathcal{A}$ .*

**Acknowledgments.** I would like to thank Professor Antonio Siconolfi for giving a nice lecture on their paper [FS2] at the Fukuoka University in October 2006, and Professor Naoki Yamada for inviting me to this lecture. This lecture is very helpful to write the manuscript.

## References

- [BCD] M. BARDI AND I. CAPUZZO-DOLCETTA, *Optimal Control and Viscosity Solutions of Hamilton-Jacobi-Bellman Equations*, Birkhäuser, Boston, 1997.
- [CI] G. CONTRERAS AND R. ITURRIAGA, *Grobal Minimizers of Autonomous Lagrangians*, 22nd Brazilian Mathematics Colloquium, Instituto de Matematica Pura e Aplicada, Rio de Janeiro, 1999.
- [CP] G. CONTRERAS AND G. PATERNAIN, *Connecting orbits between static classes for generic Lagrangian systems*, *Topology* 41 (2002), pp. 645–666.
- [DS] A. DAVINI AND A. SICONOLFI, *A generalized dynamical approach to the large time behavior of Hamilton-Jacobi equations*, *SIAM J. Math. Anal.* 38 (2006), pp. 478–502.
- [F1] A. FATHI, *Théoreme KAM et théorie de Mather pour les systèmes lagrangiens*, *C.R. Acad. Sci. Paris Sér. I* 324 (1997), pp. 1034–1046.
- [F2] A. FATHI, *Sur la convergence du semi-group de Lax-Oleinik*, *C.R. Acad. Sci. Paris Sér. I* 327 (1998), pp. 267–270.
- [F3] A. FATHI, *Weak KAM theorem in Lagrangian dynamics*, to appear.
- [FFR] A. FATHI, A. FIGALLI AND L. RIFFORD, *On the Hausdorff dimension of the Mather quotient*, to appear in *Comm. Pure Appl. Math.*
- [FS1] A. FATHI AND A. SICONOLFI, *Existence of  $C^1$  critical subsolutions of the Hamilton-Jacobi equation*, *Invent. Math.* 155 (2004), pp. 363–388.
- [FS2] A. FATHI AND A. SICONOLFI, *PDE aspects of Aubry-Mather theory for quasi-convex Hamiltonians*, *Calculus of Variations and Partial Differential Equations* 22 (2005), pp. 185–228.
- [Fu] Y. FUJITA, *A relation between uniqueness sets and total disconnectedness of quotient Aubry sets*, in progress.
- [FIL] Y. FUJITA, H. ISHII AND P. LORETI, *Asymptotic solutions of Hamilton-Jacobi equations in Euclidean  $n$  space*, *Indiana Univ. Math. J.* 55 (2006), pp. 1671–1700.
- [FU] Y. FUJITA AND K. UCHIYAMA, *Asymptotic solutions with slow convergence rate of Hamilton-Jacobi equations in Euclidean  $n$  space*, *Differential and Integral Equations* 20 (2007), pp.1185–1200.

- [I] H. ISHII, *Asymptotic solutions for large time of Hamilton-Jacobi equations in Euclidean  $n$  space*, to appear in *Annales de l'IHP-Analyse non lineaire* 25 (2008), pp.231-266
- [IM] H. ISHII AND H. MITAKE, *Representation formulas for solutions of Hamilton-Jacobi equations*, *Indiana Univ. Math. J.*, 56 (2007), pp.2159-2184.
- [M1] J. N. MATHER, *Total disconnectedness of the quotient Aubry set in low dimensions*, *Comm. Pure Appl. Math.*, 56 (2003), pp.1178-1183.
- [M2] J. N. MATHER, *Examples of Aubry sets*, *Ergod. Th. & Dynami. Sys.*, 24 (2004), pp.1667-1723.
- [S] A. SORRENTINO, *On the total disconnectedness of the quotient Aubry set*, *Ergod. Th. & Dynami. Sys.*, 28 (2007), pp.267-290.
- [W] G. T. WHYBURN, *Open mappings on locally compact spaces*, *Mem. Amer. Math. Soc.*, 1950, (1950). no. 1, American Mathematical Society, Providence, R.I.